

#### Austin Cole, MD

# Medial Patellofemoral Ligament (MPFL) Reconstruction

**Post-Operative Protocol** 

## Phase I - Maximum Protection

Weeks 0-2

- Brace- locked in full extension during all ambulation (and sleep) and WBAT
  Can unlock brace to allow 0-30 degrees for unloaded range of motion only
  - Use two crutches initially with gradual progression of weight bearing to WBAT
- Limit knee flexion to 0-30 degrees for 2 weeks Goals
  - Reduce inflammation and pain
  - 0 degrees of knee extension

Exercise progression

- Quadriceps setting, emphasize VMO, use NMES as needed
- Emphasize patellofemoral mobilizations
- Passive/active knee range of motion with 30° flexion limit
- Multi-plane straight leg raising
- Open chain hip strengthening
- Gait training

#### Weeks 2 to 4:

- Brace- locked in full extension during all ambulation (and sleep) and WBAT
- Progress ROM as tolerated to 0-90 degrees Goals
  - Reduce inflammation and pain
  - Maintain 0° of knee extension

## Weeks 4 to 6:

- Brace- able to unlock during ambulation (and sleep) and WBAT
- Progress ROM as tolerated to 0-120 degrees
- Initiate bike with light resistance
- Initiate loaded flexion 0-90 degrees Goals
  - Reduce inflammation and pain
  - Maintain 0° of knee extension

## Phase II- Progressive Stretching and Early Strengthening

#### Weeks 6 to 8:

- Discontinue brace
- Progress to full ROM as tolerated

<u>Goals</u>

- Full knee extension/hyperextension
- Progress to full knee flexion ROM
- No swelling
- Normal gait pattern

#### Exercise progression

- Continue to emphasis patella mobility
- Gait training- normalize gait pattern
- Proprioception drills
- Begin unilateral closed kinetic chain program
- Step-up progression

## Phase III- Advanced Strengthening and Endurance Training

## Weeks 8 to 10:

### <u>Goals</u>

• Full knee range of motion

Exercise progression

- o Advance stationary biking program (increase intensity)
- o Introduce treadmill walking and elliptical trainer
- Gym strengthening progression

### Weeks 10 to 12:

Exercise progression

- o Lunge progression (retro, walk, and split) as indicated
- o Swimming freestyle
- Gym strengthening progression

### Phase IV- Advanced Strengthening and Running Progression

### Weeks 12 to 14:

- Administer Preliminary functional test for physician to review
- Initiate straight line jogging at 12 weeks if proper biomechanics are demonstrated <u>Exercise progression</u>
  - Basic ladder series
  - Lateral lunge progression
  - Linear jogging progression
  - Basic plyometric box progression

### Weeks 14 to 20:

- Advance training in preparation for functional testing
- Progress plyometric training from double leg to single leg activities
  Criteria to progress to Phase V
- Pass Return to Sport test at >90% (involved vs. uninvolved limb)
  - See testing protocol- perform at 16 weeks

## Phase V- Return to Sport

### Weeks 20 to 24:

## Exercise progression

- Advance ladder, hurdle, and plyometrics
- Sport specific field/court drills
- Non-contact drills

Criteria to be released for return to sport

- Follow-up examination with the physician
- Pass return to sport functional test at >90% (involved vs. uninvolved limb)
- Display symmetry and confidence in high-speed cutting, multi-plane plyometric drills, sprinting and decelerating